


ALMA observations of 99 GHz free-free and H40 α emission from star formation in the centre of NGC 253

G. J. Bendo, R. J. Beswick, M. J. D'Cruze, C. Dickinson, G. A. Fuller, T. W. B. Muxlow

Star formation can be traced either by ultraviolet-luminous stars or by supernovae.

The wavebands commonly used to trace star formation each have drawbacks:

- Ultraviolet continuum emission from young stars is strongly affected by dust extinction.
- Optical/near-infrared recombination line emission from photoionized gas is also affected by dust extinction.
- Infrared emission from dust may include emission from dust heated by older stars.
- Synchrotron emission from supernovae appears more extended than the young stellar populations.




ALMA can detect emission from photoionized gas in two forms:

- Free-free (or Bremsstrahlung) continuum emission
- Higher order recombination line emission

This emission has two advantages over other commonly-used star formation tracers:

- It directly traces young, photoionizing stars.
- It is unaffected by dust attenuation.




Peel et al. (2011, MNRAS, 416, L99)

Detection of radio/millimetre recombination line emission has been limited by sensitivity issues.

Examples of galaxies where line emission has been detected include the following:

- Arp 220
- M82
- M83
- NGC 1365 (Fornax A)
- NGC 4945
- NGC 5253

ALMA will have the capability to detect this emission in more galaxies, making it possible to analyze the line emission in larger samples of galaxies.




Arp 220 (Rodriguez-Rico et al., 2005, ApJ, 633, 198)

Detection of radio/millimetre recombination line emission has been limited by sensitivity issues.

Examples of galaxies where line emission has been detected include the following:

- Arp 220
- M82
- M83
- NGC 1365 (Fornax A)
- NGC 4945
- NGC 5253

ALMA will have the capability to detect this emission in more galaxies, making it possible to analyze the line emission in larger samples of galaxies.




M82 (Rodriguez-Rico et al., 2004, ApJ, 616, 783)

Detection of radio/millimetre recombination line emission has been limited by sensitivity issues.

Examples of galaxies where line emission has been detected include the following:

- Arp 220
- M82
- M83
- NGC 1365 (Fornax A)
- NGC 4945
- NGC 5253

ALMA will have the capability to detect this emission in more galaxies, making it possible to analyze the line emission in larger samples of galaxies.



M83 (Zhao et al., 1996, ApJ, 472, 54)


NGC 253 is a nearby spiral galaxy with starburst activity in its nucleus.

Multiple papers dating back to 1977 had reported radio recombination line emission from the nucleus of the galaxy.

ALMA data have been published in the following papers:

- Bolatto et al. (2013, *Nature*, 499, 450)
- Leroy et al. (2014, arXiv: 1411.2836)
- Meier et al. (2015, arXiv: 1501.05694)

However, most of the focus has been on molecular gas rather than photoionized gas.




Bolatto et al. (2013, *Nature*, 499, 450)

Declination (J2000)

-25:17:10

-25:17:20



00:47:34


00:47:33

Right Ascension (J2000)

Declination (J2000)

-25:17:10

-25:17:20



00:47:34


00:47:33

Right Ascension (J2000)

Declination (J2000)

-25:17:10

-25:17:20





$v(\text{H40}\alpha)$





00:47:34

00:47:33

Right Ascension (J2000)






To use the continuum emission as a measure of free-free emission, we need to remove contributions from other sources.

Rodríguez-Rico et al. (2006, ApJ, 644, 914) and Peel et al. (2011, MNRAS, 416, L99) provide different estimates based on using data measured in different apertures and applying different techniques.

We re-fit the Rodríguez-Rico et al. data for the central 30'' without fixing the slope of the synchrotron emission.

We estimate that $70 \pm 10\%$ of the emission at 99.02 GHz is from free-free emission.




Rodríguez-Rico et al. (2006, ApJ, 644, 914)

To use the continuum emission as a measure of free-free emission, we need to remove contributions from other sources.

Rodríguez-Rico et al. (2006, ApJ, 644, 914) and Peel et al. (2011, MNRAS, 416, L99) provide different estimates based on using data measured in different apertures and applying different techniques.

We re-fit the Rodríguez-Rico et al. data for the central 30'' without fixing the slope of the synchrotron emission.

We estimate that $70 \pm 10\%$ of the emission at 99.02 GHz is from free-free emission.




Peel et al. (2011, MNRAS, 416, L99)

To use the continuum emission as a measure of free-free emission, we need to remove contributions from other sources.

Rodríguez-Rico et al. (2006, ApJ, 644, 914) and Peel et al. (2011, MNRAS, 416, L99) provide different estimates based on using data measured in different apertures and applying different techniques.

We re-fit the Rodríguez-Rico et al. data for the central 30'' without fixing the slope of the synchrotron emission.

We estimate that $70 \pm 10\%$ of the emission at 99.02 GHz is from free-free emission.




The ratio of recombination line emission to free-free emission can be used to measure the electron temperature (T_e) of the gas.

We first multiplied the continuum measurements by 0.70 to account for other sources of emission.

T_e (without adjustment):
5800-7100 K

T_e (with 0.70 adjustment):
3700-4500 K

T_e (with 0.50 adjustment):
2500-3000 K




The ratio of recombination line emission to free-free emission can be used to measure the electron temperature (T_e) of the gas.

We first multiplied the continuum measurements by 0.70 to account for other sources of emission.

T_e (without adjustment):
5800-7100 K

T_e (with 0.70 adjustment):
3700-4500 K


T_e (with 0.50 adjustment):
2500-3000 K




Paladini et al. (2004, MNRAS, 347, 237)

Both the free-free and recombination line emission can be converted to either photoionizing photon production rates (Q) or star formation rates (SFR).


We compared these rates to others published in the literature, which show significant scatter.



ALMA
 $\text{SFR} = 1.59 \pm 0.16 M_{\odot} \text{ yr}^{-1}$
 $Q = (3.0 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




ALMA
 $\text{SFR} = 1.87 \pm 0.18 M_{\odot} \text{ yr}^{-1}$
 $Q = (3.5 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




Puxley et al. (1997, ApJ, 485, 143)
 $Q = (7.0 \pm 1.5) \times 10^{53} \text{ yr}^{-1}$ (adjusted)
 low S/N

Both the free-free and recombination line emission can be converted to either photoionizing photon production rates (Q) or SFRs.

We compared these rates to others published in the literature (which show significant scatter).



ALMA
SFR = $1.59 \pm 0.16 M_{\odot} \text{ yr}^{-1}$
Q = $(3.0 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




ALMA
SFR = $1.87 \pm 0.18 M_{\odot} \text{ yr}^{-1}$
Q = $(3.5 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$

1.2 cm continuum
(no picture shown)


Ott et al. (2005, ApJ, 629, 767)
SFR = $4.9 \pm 0.5 M_{\odot} \text{ yr}^{-1}$ (adjusted)
poor spectral decomposition

Both the free-free and recombination line emission can be converted to either photoionizing photon production rates (Q) or SFRs.


We compared these rates to others published in the literature (which show significant scatter).



ALMA
SFR = $1.59 \pm 0.16 M_{\odot} \text{ yr}^{-1}$
Q = $(3.0 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




ALMA
SFR = $1.87 \pm 0.18 M_{\odot} \text{ yr}^{-1}$
Q = $(3.5 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




Rodriguez-Rico et al. (2006, ApJ, 644, 194)
Q = $1.3 \times 10^{53} \text{ yr}^{-1}$ (adjusted)
poor spectral decomposition

Both the free-free and recombination line emission can be converted to either photoionizing photon production rates (Q) or SFRs.


We compared these rates to others published in the literature (which show significant scatter).



ALMA
 $\text{SFR} = 1.59 \pm 0.16 M_{\odot} \text{ yr}^{-1}$
 $Q = (3.0 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




ALMA
 $\text{SFR} = 1.87 \pm 0.18 M_{\odot} \text{ yr}^{-1}$
 $Q = (3.5 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




Rodriguez-Rico et al. (2006, ApJ, 644, 194)
 $Q = 1.1 \times 10^{53} \text{ yr}^{-1}$ (adjusted)
 poor sensitivity or low optical depth regions only

Both the free-free and recombination line emission can be converted to either photoionizing photon production rates (Q) or SFRs.


We compared these rates to others published in the literature (which show significant scatter).



ALMA
SFR = $1.59 \pm 0.16 M_{\odot} \text{ yr}^{-1}$
Q = $(3.0 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




ALMA
SFR = $1.87 \pm 0.18 M_{\odot} \text{ yr}^{-1}$
Q = $(3.5 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




Kepley et al. (2011, ApJ, 644, 194)
Q = $1 \times 10^{53} \text{ yr}^{-1}$
poor sensitivity or low optical depth
regions only

Both the free-free and recombination line emission can be converted to either photoionizing photon production rates (Q) or SFRs.


We compared these rates to others published in the literature (which show significant scatter).



ALMA
 $\text{SFR} = 1.59 \pm 0.16 M_{\odot} \text{ yr}^{-1}$
 $Q = (3.0 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$



ALMA
 $\text{SFR} = 1.87 \pm 0.18 M_{\odot} \text{ yr}^{-1}$
 $Q = (3.5 \pm 0.3) \times 10^{53} \text{ yr}^{-1}$




Rampadarath et al. (2014, AJ, 147, 5)
 $\text{SFR} < 4.9 M_{\odot} \text{ yr}^{-1}$
 reached limits of methodology

The central starburst in NGC 253 is heavily obscured in optical wavelengths, but near-infrared recombination lines (Pa β , Br γ) have been detected (Engelbracht et al., 1998, ApJ, 505, 639).

We can compare the H40 α emission to the near-infrared lines to measure dust attenuation.


We measure attenuations that are ~ 3 dex higher than measured by Engelbracht et al.



Engelbracht et al. (1998, ApJ, 505, 639)

$$A_J = 2.00 \pm 0.36$$

$$A_K = 0.87 \pm 0.16$$



ALMA

$$A_J = 4.9 \pm 0.2$$

$$A_K = 4.2 \pm 0.2$$

Conclusions

- We detect both free-free and H40 α emission from a three-lobed structure in the centre of NGC 253.
- The T_e measured in ALMA data are consistent with previous measurements in NGC 253 as well as with measurements in the Milky Way.
- The SFR measured with ALMA data fall within the broad range of SFR values reported in the literature, but the ALMA data should be better constrained.
- Combining the ALMA line measurements with near-infrared data, we measure dust attenuation that is ~ 3 dex higher than what was determined from the near-infrared data alone.

Future Work

- ALMA free-free and recombination line emission from other galaxies could be used to cross-calibrate other star formation tracers (e.g. infrared, ultraviolet + infrared, H α + infrared emission).
- Broad recombination line measurements could be used to identify the presence of AGN, including in situations where the AGN may be heavily obscured (e.g. LIRGs).
- Line emission from composite AGN/starburst objects can be used to identify the relative contributions of these components to the overall emission from these galaxies.